

Characterizations of the Generators of Positive Semigroups on C^* - and von Neumann Algebras[†]

Jörg Rheinländer
 Institut für Mathematische Stochastik
 Georg-August-Universität
 Lotzestr. 13, D-37083 Göttingen, Germany.
 e-mail: joerg.rheinlaender@gmx.net

February 7, 2008

Abstract

Generators of positive C_0 -semigroups on C^* -algebras and C_0^* -semigroups on von Neumann algebras are examined. A characterization due to Bratteli and Robinson in the C_0 -case is proven in the C_0^* -case. Under the additional assumptions of unitality and contractivity of the semigroup another characterization of the generator is given. This result is restated for the dual and predual semigroup.

Keywords: C^* -algebra, von Neumann algebra, positivity, C_0 -semigroup, C_0^* -semigroup, infinitesimal generator

Mathematical Subject Classification (2000): 46L57, 47D06, 82C10

1 Introduction

In this note we examine infinitesimal generators of positive one parameter semigroups and of positive, unital and contractive one-parameter semigroups T_t on C^* - and von Neumann algebras. Unitality of T_t means $T_t(\mathbf{1}) = \mathbf{1}$, where $\mathbf{1}$ is the unit of the algebra.

Uniformly continuous, positive semigroups were studied by Evans and Hanché-Olsen [11]. They characterized the generators L of such semigroups by the inequalities

$$\begin{aligned} L(a^2) + aL(\mathbf{1})a &\geq L(a)a + aL(a), & \text{for all self-adjoint } a \in \mathcal{A}, \\ L(\mathbf{1}) + u^*L(\mathbf{1})u &\geq L(u^*)u + u^*L(u), & \text{for all unitary } u \in \mathcal{A}. \end{aligned}$$

[†]Part of this work is contained in the author's thesis [20].

This result was generalized by Bratteli and Robinson [2] to positive C_0 -semigroups on C^* -algebras. They derived the above inequalities with the generator L replaced by the resolvent $(\lambda - L)^{-1}$. In section 2 this result is restated for C_0^* -semigroups on von Neumann algebras. The proof of this characterization is mainly based on the proof of Bratteli and Robinson in the C_0 -case. The modification of the proof holds for both C_0 -semigroups on C^* -algebras and C_0^* -semigroups acting on von Neumann algebras.

In section 3 it is additionally assumed that the semigroups under consideration are contractive and unital. A characterization of the generators themselves can be given, which is not only a characterization in terms of their resolvents. This result for positive, unital semigroups of contractions admits a reformulation for the related dual and predual semigroups. This leads to a generalization of a result due to Kossakowski [14].

With the help of the results in section 3 it is proven in [15] that the generators of uniformly continuous, unital and positive semigroups of contractions on the C^* -algebra $\mathcal{M}(n, \mathbb{C})$ of $n \times n$ matrices are essentially the difference of two generators of completely positive semigroups.

Semigroups which satisfy this stronger notion of positivity are the natural candidates for a description of a Markovian irreversible time evolution in quantum statistical mechanics, in particular for so called reduced dynamics. These dynamics arise from a Hamiltonian dynamic through an averaging procedure over a subsystem or some internal degrees of freedom, i.e. one is interested in the family of maps $N \circ \alpha_t$, where α_t is a one-parameter group of $*$ -automorphisms of the algebra $\mathcal{A}$ and $N : \mathcal{A} \rightarrow \mathcal{B}$ is the conditional expectation from $\mathcal{A}$ into a C^* -subalgebra $\mathcal{B}$. If N and α_t are completely positive, so is the composition. Of course, $N \circ \alpha_t$ is in general not a one-parameter semigroup because of memory effects. In order to eliminate these memory effects, one usually carries out a limiting procedure, for example a weak or singular coupling limit [8]. For a detailed discussion see [10].

When the completely positive, unital semigroup is uniformly continuous, the infinitesimal generator can be written in the so-called Lindblad form [12], [16], [7], see also [4] p. 222-226. In the case of non-uniformly continuous semigroups, a canonical form of the generator is not yet known, but there are interesting results in this direction, see [13], [6] and references therein.

The description of irreversible dynamics by completely positive semigroups is not without criticism, see e.g. [4], [17] and [19].

For a detailed discussion on positive and completely positive semigroups see [15].

2 Positive semigroups

In this section we discuss positive semigroups and their generators. We state a characterization of the generators of both positive C_0 -semigroups on a C^* -

algebra and positive C_0^* -semigroups on a von Neumann algebra $\mathcal{A}$, in the C_0 -case due to Bratteli and Robinson [2].

Let us recall some definitions. If X is a Banach space then a one parameter family T_t , $t \geq 0$, of bounded linear operators on X is defined to be a strongly continuous or C_0 -semigroup if it satisfies

1. $T_t T_s = T_{t+s}$, $\forall t, s \geq 0$,
2. $T_0 = id_X$,
3. $\lim_{t \downarrow 0} \|T_t(x) - x\| = 0$, $\forall x \in X$.

Similarly, if X is a Banach space with predual space X_* a one parameter family T_t , $t \geq 0$, is defined to be a C_0^* -semigroup if it satisfies 1 and 2 from above and the weak*-continuity conditions

4. $t \mapsto \eta(T_t(x))$ is continuous $\forall x \in X$, $\forall \eta \in X_*$,
5. $x \mapsto \eta(T_t(x))$ is continuous $\forall \eta \in X_*$, $\forall t \geq 0$.

If one replaces the predual space X_* in the above definition by the dual space X^* one gets a weakly continuous semigroup. It can be proven that each weakly continuous semigroup is a C_0 -semigroup, see [23] p. 233. The generator of a semigroup T_t is defined as the linear operator L whose domain $D(L)$ consists of those $x \in X$ for which there exists a $y \in X$ such that

$$\lim_{t \downarrow 0} \frac{1}{t}(T_t(x) - x) = y = 0$$

and the action of L is then defined by $L(x) = y$. The limit is taken in the weak topology if T_t is a C_0 -semigroup and in the weak*-topology in the C_0^* -case.

We are interested in C_0 -semigroups on C^* -algebras and C_0^* -semigroups on von Neumann algebras with further properties. First we look at positive semigroups T_t , i.e. semigroups which leave the positive cone $\mathcal{A}^+$ of the algebra invariant: $T_t(\mathcal{A}^+) \subseteq \mathcal{A}^+$. In Section 3 we assume additionally the unitality $T_t(\mathbf{1}) = \mathbf{1}$ and the contractivity of the semigroups. We always assume that the C^* -algebras possess a unit $\mathbf{1}$.

In order to handle C_0 - and C_0^* -semigroups simultaneously, we use the notation $\sigma(\mathcal{A}, F)$ -continuous semigroup, where $\sigma(\mathcal{A}, F)$ is either the locally-convex topology on the C^* -algebra $\mathcal{A}$ induced by the functionals in the dual $F = \mathcal{A}^*$ corresponding to C_0 -semigroups. On the other hand $\mathcal{A}$ can be viewed as a von Neumann algebra with predual space $F = \mathcal{A}_*$. In this case $\sigma(\mathcal{A}, F)$ is the weak*-topology corresponding to C_0^* -semigroups, see [3] Section 2.5.3 and Section 3.1.2. For a discussion of the different continuity properties on C^* - and von Neumann algebras, see [3] Section 3.2.2.

A linear mapping $T : D(T) \rightarrow \mathcal{A}$ on a C^* - and von Neumann algebra $\mathcal{A}$ respectively with domain $D(T) \subseteq \mathcal{A}$ is called symmetric, if $T(x^*) = T(x)^*$ for all $x \in D(T)$.

For the theory of C^* - and von Neumann algebras the reader is referred to [3] and [22]. References for the theory of one-parameter semigroups are e.g. [3], [9], [18]. In [3] the theories of C_0 - and C_0^* -semigroups are developed together.

Theorem 1 *Let L be the generator of a $\sigma(\mathcal{A}, F)$ -continuous semigroup T_t of symmetric operators on a unital algebra $\mathcal{A}$. Then the following conditions are equivalent:*

1. T_t is positive for all $t \geq 0$.
2. $(\lambda - L)^{-1}$ is positive for all large $\lambda > 0$.
3. $(\lambda - L)^{-1}(a^2) + a(\lambda - L)^{-1}(\mathbf{1})a \geq (\lambda - L)^{-1}(a)a + a(\lambda - L)^{-1}(a)$, for all self-adjoint $a \in \mathcal{A}$ and all large $\lambda > 0$.
4. $(\lambda - L)^{-1}(\mathbf{1}) + u^*(\lambda - L)^{-1}(\mathbf{1})u \geq (\lambda - L)^{-1}(u^*)u + u^*(\lambda - L)^{-1}(u)$, for all unitary $u \in \mathcal{A}$ and all large $\lambda > 0$.
5. $T_t(a^2) + aT_t(\mathbf{1})a \geq T_t(a)a + aT_t(a)$, for all self-adjoint $a \in \mathcal{A}$ and $t \geq 0$.
6. $T_t(\mathbf{1}) + u^*T_t(\mathbf{1})u \geq T_t(u^*)u + u^*T_t(u)$, for all unitary $u \in \mathcal{A}$ and $t \geq 0$.

Remark: The proof of the C_0^* -case requires a modification of the proof of the C_0 -case in [2]. There one uses a convergence argument to prove the equivalence of 5. and 6. to 1. – 4. This argument does not work in the case of C_0^* -semigroups. In order to overcome this difficulty a Laplace transform argument is used in the following proof. For the sake of convenience, the complete proof is given.

Proof: 1. $\Rightarrow$ 2.: This follows from the Laplace transformation:

$$(\lambda - L)^{-1}(a) = \int_0^\infty e^{-\lambda t} T_t(a) dt, \quad \text{for all } \lambda > 0, \quad \text{for all } a \in \mathcal{A}.$$

The usual relation for C_0 -semigroups between the resolvent and the Laplace transform holds also for C_0^* -semigroups, see [3] Proposition 3.1.6.

2. $\Rightarrow$ 1.:

$$T_t(a) = \lim_{n \rightarrow \infty} \left(\frac{n}{t} \left(\frac{n}{t} - L \right)^{-1} \right)^n(a), \quad \text{for all } t > 0, \quad \text{for all } a \in \mathcal{A},$$

where the limit is taken in the $\sigma(\mathcal{A}, F)$ -topology. Thus, the semigroup is positive whenever the resolvent is positive for λ sufficiently large.

1. $\Rightarrow$ 3. (2. $\Rightarrow$ 4.): In the case of uniformly continuous semigroups and bounded generators respectively conditions 1. and 2. are equivalent to one of the following two conditions [11]:

$$L(a^2) + aL(\mathbf{1})a \geq L(a)a + aL(a), \quad \text{for all self-adjoint } a \in \mathcal{A}, \quad (1)$$

$$L(\mathbf{1}) + u^*L(\mathbf{1})u \geq L(u^*)u + u^*L(u), \quad \text{for all unitary } u \in \mathcal{A}. \quad (2)$$

With this result in mind, it is sufficient to show the equivalence of 1. and 2. to the following condition.

7. *The uniformly continuous semigroup S_t with generator $(\lambda - L)^{-1}$ is positive for all $t \geq 0$ and large $\lambda > 0$.*

2. $\Rightarrow$ 7.: Follows from

$$S_t = e^{t(\lambda - L)^{-1}} = \sum_{n=0}^{\infty} \frac{t^n}{n!} (\lambda - L)^{-n}.$$

7. $\Rightarrow$ 1.: Let U_t^λ be the uniformly continuous semigroup whose generator L_λ is given by the Yosida approximation of L , i.e. $L_\lambda := \lambda L(\lambda - L)^{-1} = \lambda^2(\lambda - L)^{-1} - \lambda$. U_t^λ is positive if S_t is positive, since

$$U_t^\lambda = e^{t(\lambda^2(\lambda - L)^{-1} - \lambda)} = e^{-t\lambda} e^{\lambda^2 t(\lambda - L)^{-1}} = e^{-t\lambda} S_{\lambda^2 t}.$$

T_t is the $\sigma(\mathcal{A}, F)$ -limit of U_t^λ for $\lambda \rightarrow \infty$, see [18] Theorem 1.5.5. for the result in the C_0 -case the C_0^* -result follows by duality. So T_t is positive.

5. $\Rightarrow$ 3. (6. $\Rightarrow$ 4.): We have

$$T_t(a^2) + aT_t(\mathbf{1})a - (T_t(a)a + aT_t(a)) \geq 0,$$

for all $t \geq 0$ and all self-adjoint (unitary $a \in \mathcal{A}$). So

$$\begin{aligned} & (\lambda - L)^{-1}(a^2) + a(\lambda - L)^{-1}(\mathbf{1})a - ((\lambda - L)^{-1}(a)a + a(\lambda - L)^{-1}(a)) \\ &= \int_0^\infty e^{-\lambda t} (T_t(a^2) + aT_t(\mathbf{1})a - (T_t(a)a + aT_t(a))) dt \geq 0 \end{aligned}$$

follows by Laplace transformation.

1. $\Rightarrow$ 5. (6.): Provided $T_t \geq 0$ for all $t \geq 0$,

$$e^{sT_t} = \sum_{n=0}^{\infty} \frac{(sT_t)^n}{n!}$$

is positive for all $s \geq 0$ and $t \geq 0$. Thus, the estimates (1) and (2) hold for the generator T_t of the semigroup e^{sT_t} .

3 Positive, unital semigroups of contractions

In the previous section we discussed $\sigma(\mathcal{A}, F)$ -continuous, positive semigroups. Now we assume in addition that the semigroup is unital and contractive. These properties allow a characterization of the generator which is different to the characterizations in Theorem 1. There the characterizations are in terms of the resolvent of the generator, in the following we provide a characterization in terms of the generator only.

Theorem 2 *Let L be the generator with domain $D(L)$ of a $\sigma(\mathcal{A}, F)$ -continuous semigroup T_t of contractions on $\mathcal{A}$. The following two conditions are equivalent:*

1. T_t is positive and unital.
2. $\mathbf{1} \in D(L)$, $L(\mathbf{1}) = 0$ and L is symmetric.

Proof: 1. $\Rightarrow$ 2.: $\mathbf{1} \in D(L)$ and $L(\mathbf{1}) = 0$ follow from $\lim_{h \downarrow 0} \frac{1}{h}(T_h(\mathbf{1}) - \mathbf{1}) = 0$, where the limit is taken in the $\sigma(\mathcal{A}, F)$ -topology.

Every element $x \in \mathcal{A}$ can be decomposed into a linear combination of four positive elements $a, b, c, d \in \mathcal{A}^+$ such that

$$x = a - b + i(c - d).$$

Since T_t is positive and linear we get with the help of this decomposition

$$T_t(x^*) = T_t(x)^*,$$

thus T_t is symmetric. Let $x \in D(L)$, then by using the uniform-continuity of the $*$ -map

$$\begin{aligned} L(x)^* &= \lim_{h \rightarrow \infty} \frac{1}{h}(T_h(x) - x)^* \\ &= \lim_{h \rightarrow \infty} \frac{1}{h}(T_h(x^*) - x^*), \end{aligned}$$

the limit is performed in the $\sigma(\mathcal{A}, F)$ -topology. Thus $x^* \in D(L)$ and $L(x^*) = L(x)^*$.

2. $\Rightarrow$ 1.: Property 2. reads in terms of the resolvent $(\lambda - L)^{-1}(\mathbf{1}) = \frac{1}{\lambda}\mathbf{1}$. By approximation in the $\sigma(\mathcal{A}, F)$ -topology we get

$$T_t(\mathbf{1}) = \lim_{n \rightarrow \infty} \left(\frac{n}{t} \left(\frac{n}{t} - L\right)^{-1}\right)^n(\mathbf{1}) = \lim_{n \rightarrow \infty} \left(\frac{n}{t}\right)^n \left(\frac{t}{n}\right)^n(\mathbf{1}) = \mathbf{1},$$

which proves the unitality of T_t .

By assumption L is the generator of a semigroup of contractions. Therefore the range of $\lambda - L$ is the whole algebra $\mathcal{A}$ for $\lambda > 0$. For $x \in \mathcal{A}$ exists a unique

element $y \in D(L)$, such that $(\lambda - L)^{-1}(x) = y$. Clearly, $\lambda - L$ is symmetric for $\lambda > 0$. Therefore

$$\begin{aligned} ((\lambda - L)^{-1}(x))^* &= ((\lambda - L)^{-1}(\lambda - L))(y^*) \\ &= (\lambda - L)^{-1}(x^*). \end{aligned}$$

The semigroup T_t can be defined by $T_t(x) = \lim_{n \rightarrow \infty} (\frac{n}{t}(\frac{n}{t} - L)^{-1})^n(x)$. In the C_0^* -case the limit is taken in the weak- $*$ -topology on the von Neumann algebra $\mathcal{A}$. The weak- $*$ -topology on $\mathcal{A}$ is the relative σ -weak topology on $\mathcal{A}$ and the involution $*$ is continuous in this topology. So one can interchange the involution and the limit, thus T_t is symmetric. In the C_0 -case the limit is taken in the weak topology on the C^* -algebra $\mathcal{A}$. Every weakly continuous semigroup is also strongly continuous, see [23] p.233, and the involution $*$ is continuous with respect to the uniform topology on the C^* -algebra $\mathcal{A}$. As in the von Neumann case we can interchange the limit with the involution and the symmetry of T_t follows.

Now let $a \in \mathcal{A}^+$ and $a_1 := \frac{a}{\|a\|}$. $\mathcal{A}^+$ is a convex cone, thus $a_1 \in \mathcal{A}^+$ and $\|a_1\| = 1$. So the spectrum, $\sigma(a_1)$, of a_1 is contained in $[0, 1]$. Therefore

$$1 - \sigma(a_1) = \sigma(\mathbf{1} - a_1) \subseteq [0, 1]$$

and

$$\|\mathbf{1} - a_1\| \leq 1.$$

Thus

$$\begin{aligned} \|\mathbf{1} - T_t(a_1)\| &= \|T_t(\mathbf{1}) - T_t(a_1)\| \\ &\leq \|T_t\| \|\mathbf{1} - a_1\| \\ &\leq 1. \end{aligned}$$

The semigroup T_t is symmetric for all $t \geq 0$, hence $T_t(a_1) \in \mathcal{A}^{sa}$. With this

$$1 - \sigma(T_t(a_1)) = \sigma(\mathbf{1} - T_t(a_1)) \subseteq [-1, 1].$$

It follows that $\sigma(T_t(a_1)) \subseteq [0, 2]$, therefore $T_t(a_1) \in \mathcal{A}^+$. One concludes

$$T_t(a) = \|a\| T_t(a_1) \in \mathcal{A}^+,$$

thus T_t is positive, which proves the theorem.

Remark: The direction 2. $\Rightarrow$ 1. can be sharpened using the fact, that a unital, contractive map between two unital C^* -algebras and two von Neumann algebras respectively is positive, see [21], [3] Corollary 3.2.6.

Theorem 3 *Let L be the generator of a $\sigma(\mathcal{A}, F)$ -continuous semigroup T_t of contractions on $\mathcal{A}$.*

$$\mathbf{1} \in D(L), \quad L(\mathbf{1}) = 0 \quad \Rightarrow \quad T_t \text{ is positive.}$$

Proof: The unitality is proven as in Theorem 2. Since T_t is a contraction, i.e. $\|T_t\| \leq 1$ for all $t \geq 0$, we have $\|T_t\| = 1$. The conclusion now follows from the above remark.

Corollary 1 *A unital $\sigma(\mathcal{A}, F)$ -continuous semigroup of contractions on an operator algebra is positive.*

In the following, the above characterization is restated for the dual and the predual semigroup respectively.

Let T_t be a C_0 -semigroup of contractions on a C^* -algebra $\mathcal{A}$. Then, the adjoint operators $T_t^* : \mathcal{A}^* \rightarrow \mathcal{A}^*$ form a C_0^* -semigroup of contractions on the dual space $\mathcal{A}^*$ of $\mathcal{A}$, see [5]. The generator of T_t^* is given by the adjoint operator L^* of the generator L of the semigroup T_t . In $\mathcal{A}^*$, the set

$$\mathcal{A}^{*+} := \{\eta \in \mathcal{A}^* : \eta(a) \geq 0 \ \forall a \in \mathcal{A}^+\}$$

is a positive cone [1]. Let $\eta \in \mathcal{A}^{*+}$ and T_t be a positive C_0 -semigroup. Then $\eta(T_t(a)) \geq 0$ for all $a \in \mathcal{A}^+$. So T_t^* is also positive since $(T_t^*(\eta))(a) = \eta(T_t(a))$. For a unital semigroup T_t , one obtains $(T_t^*(\eta))(1) = \eta(1)$ for all $\eta \in \mathcal{A}^*$. $L(1) = 0$ implies $L^*(\eta)(1) = 0$ for all $\eta \in \mathcal{A}$.

In the case that T_t is a C_0^* -semigroup of contractions on a von Neumann algebra $\mathcal{A}$, there exists a C_0 -semigroup T_t^* of contractions on the predual $\mathcal{A}_*$ which is adjoint to T_t . The unitality and positivity of the semigroup T_t and $L(1) = 0$ can be transferred to T_t^* and L^* as in the case of a C^* -algebra. Finally, one gets

Corollary 2 *Let T_t^* be a C_0^* -semigroup of contractions on $\mathcal{A}^*$ the dual of a C^* -algebra $\mathcal{A}$ or a C_0 -semigroup of contractions on the predual $\mathcal{A}_*$ of a von Neumann algebra $\mathcal{A}$. Denote the generator of T_t^* by L^* . The following two conditions are equivalent:*

1. T_t^* is positive and $T_t^*(\eta)(1) = \eta(1)$ for all $\eta \in \mathcal{A}^*$ or for all $\eta \in \mathcal{A}_*$.
2. $L^*(\eta)(1) = 0$ for all $\eta \in D(L^*)$.

If one applies the last result to $\mathcal{A}_* = L^1(\mathcal{H})$ the Banach space of trace class operators and restricts the semigroup T_t^* to the real Banach space $L^1(\mathcal{H})^{sa}$ of all selfadjoint operators in $L^1(\mathcal{H})$, one sees that the result due to Kossakowski [14] is contained in Corollary 2.

Acknowledgements: The author would like to thank H. Hering, H. Leschke and M. Schütz. This work was partially supported by the Deutsche Forschungsgemeinschaft under grant no. Le 330/12-1.

References

- [1] Batty, C. J. K., Robinson, D. W.: Positive one-parameter semigroups on ordered Banach spaces, *Acta Appl. Math.* **1**, 221-296 (1984)

- [2] Bratteli, O., Robinson, D. W.: Positive C_0 -semigroups on C^* -algebras, *Math. Scand.* **49**, 259-274 (1981)
- [3] Bratteli, O., Robinson, D. W.: *Operator Algebras and Quantum Statistical Mechanics 1, sec. ed.*, Springer Verlag, Berlin (1987)
- [4] Bratteli, O., Robinson, D. W.: *Operator Algebras and Quantum Statistical Mechanics 2, sec. ed.*, Springer Verlag, Berlin (1996)
- [5] Butzer, P. L., Berens, H.: *Semi-Groups of Operators and Approximation*, Springer Verlag, Berlin (1967)
- [6] Chebotarev, A. M., Fagnola, F.: Sufficient conditions for conservativity of quantum dynamical semigroups, *J. Funct. Anal.* **118**, 131-153 (1993)
- [7] Christensen, E., Evans, D. E.: Cohomology of operator algebras and quantum dynamical semigroups, *J. London Math. Soc. (2)* **20**, 358 (1979)
- [8] Dümcke, R., Spohn, H.: The proper form of the generator in the weak coupling limit, *Z. Phys.* **B34**, 419-422 (1979)
- [9] Engel, K.-J., Nagel, R.: *One-Parameter Semigroups for Linear Evolution Equations*, Springer Verlag, Berlin (2000)
- [10] Evans, D. E.: Generators of dynamical semigroups, *Operator algebras and applications, Proc. Symp. Pure Math. 38, Part 2, Kingston/Ont. 1980*, 377-379 (1982)
- [11] Evans, D. E., Hanche-Olsen, H.: The generators of positive semigroups, *J. Funct. Anal.* **32**, 207-212 (1979)
- [12] Gorini, V., Kossakowski, A., Sudarshan, E. C. G.: Completely positive dynamical semigroups of N-level systems, *J. Math. Phys.* **17**, 821-825 (1976)
- [13] Holevo, A. S.: On the structure of covariant dynamical semigroups, *J. Funct. Anal.* **131**, 255-278 (1995)
- [14] Kossakowski, A.: On necessary and sufficient conditions for the generators of a quantum dynamical semi-group, *Bull. Acad. Polon. Sci., Sér. Sci. Math. Astronom. Phys.* **20**, 1021-1025 (1972)
- [15] Leschke, H., Rheinländer, J., Schütz, M.: Generators of positive and completely positive dynamical semigroups, Preprint, Erlangen (2001)
- [16] Lindblad, G.: On the generators of quantum dynamical semigroups, *Commun. Math. Phys.* **48**, 119-130 (1976)
- [17] Majewski, W. A.: Dynamical semigroups in the algebraic formulation of statistical mechanics, *Fortschr. Phys.* **32**, 89-133 (1984)
- [18] Pazy, A.: *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer Verlag, New York (1983)

- [19] Pechukas, P.: Reduced dynamics need not to be completely positive, *Phys. Rev. Lett.* **73**, 1060-1062 (1994); Alicki, R.: Comment on ‘Reduced dynamics need not to be completely positive’, *Phys. Rev. Lett.* **75**, 3020 (1995); Pechukas, P.: Reply on ‘Comment on ‘Reduced dynamics need not to be completely positive’’, *Phys. Rev. Lett.* **75**, 3021 (1995)
- [20] Rheinländer, J.: Charakterisierung der Generatoren dynamischer Halbgruppen, *Mathematica Gottingensis* **16**, (1997)
- [21] Russo, B., Dye, H. A.: A note on unitary operators in C^* -algebras, *Duke. Math. J.* **33**, 413-416 (1966)
- [22] Takesaki, M.: *Theory of Operator Algebras 1*, Springer Verlag, Berlin (1979)
- [23] Yosida, K.: *Functional Analysis, 6th ed.*, Springer Verlag, Berlin (1980)